

MMA

205

COMPLEX ANALYSIS


TUTORIALS

(TUT-2)

- Aryaman Maithani

General Info

- ① Keep yourself muted unless you wish to say something.
- ② If you wish to say something, directly unmute yourself and speak. No point in using "Raise hand" function.
- ③ If you feel like I've made a mistake, just point it out there.



- ④ Alternately, you may put it in the chat and someone else may point out your/my mistake. I would view the chat later and address it if the need may be.
- ⑤ When unmuting and saying something, I won't be seeing your name since I'll be on another screen. It is encouraged (but not forced) that you just mention your name at the beginning.
(Since it's being recorded, I understand if you don't want to.)

§ TUTORIAL - 1

25.08.2020

Recap. A set $U \subset \mathbb{C}$ is open if for every $z_0 \in U$, $\exists \delta > 0$ s.t.

$$B_\delta(\underline{z_0}) \subset U.$$

$$\{z \in \mathbb{C} : |z - z_0| < \delta\}$$


open ball of radius δ at z_0

δ -neighbourhood of z_0

- A subset $K \subset \mathbb{C}$ is said to be closed if $\mathbb{C} \setminus K$ is open.
- A subset $P \subset \mathbb{C}$ is said to be path-conn. if for every $z_0, z_1 \in P$, there exists a continuous function

$$[\sigma : [0, 1] \rightarrow P] \text{ s.t.}$$

$$\sigma(0) = z_0 \text{ and } \sigma(1) = z_1.$$



Differentiation

↗ open subset of \mathbb{C}

A function $f: \Omega \rightarrow \mathbb{C}$ is said to be (complex) differentiable at $z_0 \in \Omega$

if $\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$ exists.

↘ $f'(z_0)$, if it exists

CR \rightarrow Cauchy - Riemann Equations

$$u_x = v_y$$

$$u_y = -v_x$$

Any (complex) function $f: \Omega \rightarrow \mathbb{C}$ can be viewed as a function

$$f: \Omega \rightarrow \mathbb{R}^2.$$

↳ also regard as a subset of \mathbb{R}^2

$$f(x, y) = (u(x, y), v(x, y))$$

where $v, u: \Omega \rightarrow \mathbb{R}$. (Real-valued)

Thus, u_x, u_y, v_x, v_y make sense.

What CR eqⁿs tell us is:

If f is (complex) differentiable

at $z_0 = x_0 + iy_0$, then

$$u_x(x_0, y_0) = v_y(x_0, y_0),$$

$$u_y(x_0, y_0) = -v_x(x_0, y_0).$$

(existence of partial deriv. is part of the thm.)

One more thing that makes sense is the total derivative. (Since f is being viewed as a function from $\Omega \subseteq \mathbb{R}^2$ to \mathbb{R}^2 .)

Thm. If f is complex diff at $z_0 \in \Omega$, then the total derivative of f at z_0 exists.

Remark.

Total deriv. $\not\Rightarrow$ Complex der.

CR eqⁿs $\not\Rightarrow$ Complex der.

Q4.

Total Deriv. & CR eq's at $(x_0, y_0) \iff$ Complex diff. at $z_0 = x_0 + iy_0$.


MA105:

Let $f: \Omega \rightarrow \mathbb{R}^2$ with $f = (u, v)$.
 If u_x, u_y, v_x, v_y are continuous at some $(x_0, y_0) \in \Omega$, then f has a total derivative at (x_0, y_0) .

f is said to have total derivative $[A \rightarrow \text{linear trans. from } \mathbb{R}^2 \text{ to } \mathbb{R}^2]$ at $\bar{x}_0 = (x_0, y_0)$ if

$$\lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{R}^2}} \frac{\|f(\bar{x}_0 + h) - f(\bar{x}_0) - Ah\|}{\|h\|} = 0.$$

$f: \mathbb{R} \rightarrow \mathbb{R}$



$f: \mathbb{C} \rightarrow \mathbb{C}$

graph?
↳ 4D

4. Check for real differentiability and holomorphicity:

✓ 1. $f(z) = c \longrightarrow$

2. $f(z) = z$

3. $f(z) = z^n, n \in \mathbb{Z}$

4. $f(z) = \operatorname{Re}(z)$

✓ 5. $f(z) = |z|$

6. $f(z) = |z|^2$

✓ 7. $f(z) = \bar{z} \longrightarrow$ same as (4)

8. $f(z) = \begin{cases} \frac{z}{\bar{z}} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$

bit.ly/ca-205

Real diff everywhere & complex diff precisely at 0

1. $f(z) = c$. Take $z_0 \in \mathbb{C}$.

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{z \rightarrow z_0} \frac{0}{z - z_0} = \underline{0}.$$

Thus, f is (complex) diff everywhere.
 \hookrightarrow holo
 \hookrightarrow real diff.

3. $f(z) = z^n$.

① $n \geq 0$ f is defined on \mathbb{C} .
 (conv: $0^0 = 1$)

$$\begin{aligned} \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} &= \lim_{z \rightarrow z_0} \frac{z^n - z_0^n}{z - z_0} = \lim_{z \rightarrow z_0} \frac{(z - z_0)(z^{n-1} + z^{n-2}z_0 + \dots + z z_0^{n-2} + z_0^{n-1})}{(z - z_0)} \\ &= \lim_{z \rightarrow z_0} \left(\sum_{i=0}^{n-1} z^i z_0^{n-1-i} \right) \end{aligned}$$

$$= \sum_{j=0}^{n-1} z_0^{n-1} = n z_0^{n-1}.$$

Thus, f is (complex) diff. at each $z \in \mathbb{C}$.

Thus, f is holo. on \mathbb{C} . (defn)

Thus, f is real diff on \mathbb{C} . (Thm.)

② $n < 0$. Here f is defined on $\mathbb{C} \setminus \{0\}$.

On $\mathbb{C} \setminus \{0\}$, $f(z) = z^n$ is non-zero.

Thus, $\frac{1}{f}$ is diff at $z_0 \Leftrightarrow f$ is diff at z_0 .

This, we know by case 1.

$$\left[\frac{1}{f(z)} = z^{-n} \quad \text{and} \quad -n > 0. \right]$$

④ $f(z) = \operatorname{Re}(z)$

$$f(z) = f(x, y) = u(x, y) + i v(x, y).$$

$$\text{Here } u(x, y) = x,$$

$$v(x, y) = 0.$$

Clearly, u_x, v_y, v_x, u_y exist everywhere and are continuous.

Thus, f is real diff. everywhere.

Holomorphic?
(Complex diff.)

$$\left. \begin{array}{l} u_x(x_0, y_0) = 1 \\ u_y(x_0, y_0) = 0 \\ v_x(x_0, y_0) = 0 \\ v_y(x_0, y_0) = 0 \end{array} \right\} \text{for all } (x_0, y_0) \in \mathbb{R}^2$$

$$\rightarrow u_x(x_0, y_0) \neq v_y(x_0, y_0) \text{ for any } (x_0, y_0) \in \mathbb{R}^2$$

Thus, the CR eqⁿs hold NOWHERE.

Thus, f is not complex diff. anywhere.

Remark. f is cont. everywhere but diff. nowhere.

$$\textcircled{5} \quad f(z) = |z|$$

$$f(x, y) = \sqrt{x^2 + y^2} + i0$$

$$u(x, y) = \sqrt{x^2 + y^2}$$

$$v(x, y) = 0$$

First note: u_x, u_y don't exist at $(0, 0)$.

$$\hookrightarrow \lim_{x \rightarrow 0} \frac{u(x, 0) - u(0, 0)}{x} = \lim_{x \rightarrow 0} \frac{|x|}{x}$$

Thus: f is NOT real diff at $(0, 0)$.

On $\mathbb{R}^2 \setminus \{(0, 0)\}$: u & v have cont.
partial derivatives.
(Check!)

Thus, f is real diff on $\mathbb{R}^2 \setminus \{(0, 0)\}$.

Now, we check for holo:

$$v_x(x_0, y_0) = v_y(x_0, y_0) = 0 \quad \forall (x_0, y_0) \in \mathbb{R}^2$$

$$\mathbb{R}^2 \setminus \{(0, 0)\}: \quad u_x(x_0, y_0) = \frac{x_0}{\sqrt{x_0^2 + y_0^2}} = v_y = 0$$

$$u_y(x_0, y_0) = \frac{y_0}{\sqrt{x_0^2 + y_0^2}} = -v_x = 0$$

For CR eqⁿs to hold, we need $x_0 = 0$
& $y_0 = 0$.
 $\rightarrow \leftarrow$

To conclude: f is real diff on $\mathbb{R}^2 \setminus \{(0,0)\}$.
 f is complex diff. NOWHERE.

5. Show that the CR equations take the form

$$u_r = \frac{1}{r} v_\theta \quad \& \quad v_r = -\frac{1}{r} u_\theta$$

in polar coordinates.

Initiate slides.

$$f(r, \theta) = u(r, \theta) + i v(r, \theta). \\ f(re^{i\theta}) =$$

Consider $z_0 \neq 0$.

Thus, $z_0 = r_0 e^{i\theta_0}$ for some
 $r_0 > 0$
and $\theta_0 \in \mathbb{R}$.

Suppose f is diff at $z_0 = r_0 e^{i\theta_0}$.

Then,

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad \text{exists}$$

Compute \nearrow in 2 diff. ways.

① Fix $\theta = \theta_0$ and let $r \rightarrow r_0$.

$$f'(z_0) = \lim_{r \rightarrow r_0} \frac{f(r, \theta_0) - f(r_0, \theta_0)}{\underline{r}e^{i\theta_0} - r_0\underline{e}^{i\theta_0}}$$

$$= \underline{e^{-i\theta_0}} \lim_{r \rightarrow r_0} \left\{ \underbrace{\frac{u(r, \theta_0) - u(r_0, \theta_0)}{r - r_0}}_{\in \mathbb{R}} + i \underbrace{\left(\frac{v(r, \theta_0) - v(r_0, \theta_0)}{r - r_0} \right)}_{\in \mathbb{R}} \right\}$$

\downarrow this must exist and by defⁿ, it $u_r(r_0, \theta_0)$

$$f'(z) = e^{-i\theta_0} \left\{ u_r(r_0, \theta_0) + i v_r(r_0, \theta_0) \right\} \quad (*)$$

② Fix $r = r_0$ and let $\theta \rightarrow \theta_0$.

$$\frac{1}{r_0} \lim_{\theta \rightarrow \theta_0} \left\{ \underbrace{\frac{u(r_0, \theta) - u(r_0, \theta_0)}{e^{i\theta} - e^{i\theta_0}}}_{\text{this must exist}} + i \left(\frac{v(r_0, \theta) - v(r_0, \theta_0)}{e^{i\theta} - e^{i\theta_0}} \right) \right\}$$

$$\lim_{\theta \rightarrow \theta_0} \frac{u(r_0, \theta) - u(r_0, \theta_0)}{\theta - \theta_0} \cdot \frac{\theta - \theta_0}{e^{i\theta} - e^{i\theta_0}}$$

\downarrow this limit can be computed as $\frac{1}{ie^{i\theta_0}}$

\rightarrow this exists $\neq 0$


$$f'(z_0) = \frac{1}{r_0} \left\{ \frac{u_0(r_0, \theta_0)}{ie^{i\theta_0}} + i \frac{v_0(r_0, \theta_0)}{ie^{i\theta_0}} \right\} \quad (**)$$

Compare (*) & (**) to get the answer.

3. Show that if U is a path connected open set in \mathbb{C} , so is U minus finite set.
(Try to prove it as rigorously as you can)

Solⁿ. Note: If $w \in U$, then $U \setminus \{w\}$ is open.
(Proof?)

If I show $U \setminus \{w\}$ is path-connected, then I am done. (Why?)



(Want: $\sigma: [0,1] \rightarrow U \setminus \{w\}$ continuous
s.t. $\sigma(0) = z_0$,
 $\sigma(1) = z_1$)



Let $z_0, z_1 \in U \setminus \{w\}$.

Since U is path-connected, \exists cts. $\sigma: [0,1] \rightarrow U$
s.t. $\sigma(0) = z_0, \sigma(1) = z_1$

① $\sigma(x) \neq w$ for any $x \in [0, 1]$.

σ then actually maps into $U(w)$. Then done. (why?)

② $\sigma(x) = w$ for some $x \in [0, 1]$.



Using the fact that U is open,

I can find a closed ball

$$\bar{B} = \{ z \in \mathbb{C} : |z - w| \leq \delta \}.$$

($\exists \delta > 0$)

s.t.

- ① $z_0 \notin \bar{B}$, ② $z_1 \notin \bar{B}$, ③ $\bar{B} \subset U$.

$\delta_1 \nearrow$
happens

why does such a \bar{B} exist?


$\delta_3/2 \nearrow$

Since U is open, $\exists \delta > 0$ s.t. $B_\delta(\omega) \subset U$.

Then, $\overline{B_{\delta/2}(\omega)} \subset B_\delta(\omega) \subset U$.

Now, consider

$$C := \sigma^{-1}(\overline{B}) = \{x \in [0, 1] : \sigma(x) \in \overline{B}\}.$$



closed

Thus, C is closed.

$$[C \subset [0, 1]]$$

Why? Because σ is continuous.

[(Ex) If $f: \mathbb{R} \rightarrow \mathbb{C}$ is cts, then $f^{-1}(K)$ is closed whenever $K \subset \mathbb{C}$ is closed.]



① C is closed.

② C is bounded. (Why?)

\Downarrow

③ $\sup C$ & $\inf C$ exist

④ $s \in C$ and $t \in C$] \rightarrow this is because C is closed



Thus, the green path is
one such desired path!

① Show that $f(x)$ has $\deg f(x)$ roots.

Assume FTA.

f has one root x_0 .

Then, $f(x) = (x - x_0) g(x)$

↓
 $\deg f(x) - 1$
↓ use induc.

Note: $f(x) = 0$

$(\Rightarrow) x = x_0$ or $g(x) = 0$

Thus, the complete (multi)set of roots is
 $\{x_0\} \cup \{x_i : g(x_i) = 0\}^{n-1}$

1. A complex polynomial of degree n has exactly n roots. (Assuming fundamental theorem of algebra)

2. Show that a real polynomial that is irreducible has degree at most two. i.e., if

$$f(x) = a_0 + a_1x + \dots + a_nx^n, \quad a_i \in \mathbb{R},$$

then there are non-constant real polynomials g and h such that $f(x) = g(x)h(x)$ if $n \geq 3$.

② $f(x) \in \mathbb{R}[x]$. $\deg f(x) \geq 3$.

S.T. $f(x) = g(x)h(x)$

\hookrightarrow non-const real poly

① f has a root in \mathbb{R} . Done ✓ (why?)
 $f(x) = (x - \alpha_0)g(x)$
 \hookrightarrow non-const

② f has no root in \mathbb{R} .

(By FTA), f has a root $\alpha_0 \in \mathbb{C}$.

Then, $f(\bar{\alpha}_0) = 0$ as well. (why?)

$x - \alpha_0$ & $x - \bar{\alpha}_0$ are factors of $f(x)$

$\Rightarrow f(x) = \underbrace{(x - \alpha_0)(x - \bar{\alpha}_0)}_{\substack{\in \mathbb{R}[x] \\ \text{"} g(x) \text{"}}} h(x)$
 $\hookrightarrow \in \mathbb{C}[x]$

$f(x) = g(x)h(x)$

$f(x), g(x) \in \mathbb{R}[x]$

Thus, $h(x) \in \mathbb{R}[x]$.

$\hookrightarrow \deg = n - 2 \geq 1$.

Poly. div.

$f(x) = g(x)q(x) + r(x)$ in $\mathbb{R}[x]$

$\deg r(x) < 2$.

\hookrightarrow in $\mathbb{C}[x]$ as well

- $r(x) = 0$ since $g(x) \mid f(x)$ in $\mathbb{C}[x]$