

$$\int (\overset{\circ}{\underbrace{\quad}} \underbrace{\quad} \overset{\circ}{\underbrace{\quad}}) dx$$

MA 526

Commutative Algebra

Professor:

Notes By: Aryaman Maithani

Spring 2020-21

Last Updated: May 14, 2022

Contents

0	Notations and Conventions	1
1	Associated primes of ideals and modules	2
2	Artinian rings and Artinian modules	7
3	Integral Extensions of Rings	11
4	Dimension Theory of Affine Algebra over Fields	15
5	Graded Rings and Graded Modules	19
6	Dimension Theory of Finite Modules over Noetherian Local Rings	25

§0. Notations and Conventions

1. R will denote a commutative ring with unity. Modules will be unital.
2. If M is an R -module, then $\text{ann}_R(M) = \{r \in R : rM = 0\}$.
A similar definition holds for elements and subsets of M .
3. $\mathcal{N}(R)$ denotes the nilradical of R .
4. $\mathcal{J}(R)$ denotes the Jacobson radical of R .
5. $\text{Spec}(R)$ denotes the set of prime ideals of R .
6. $\text{mSpec}(R)$ denotes the set of maximal ideals of R .
7. $N \leq M$ is read as “ N is a submodule of M .”
8. $I \trianglelefteq R$ is read as “ I is an ideal of R .”
9. For an integral domain R , $Q(R)$ denotes its field of fractions.
10. k denotes a field. If k is algebraically closed, we write this as $k = \bar{k}$.
11. When we say that “ M is a finite R -module,” we mean that “ M is a finitely generated R -module.”
12. When we write “ $P(n)$ is true for $n \gg 0$,” we mean that “there exists $N \in \mathbb{N}$ such that $P(n)$ is true for all $n \geq N$.” A similar definition holds for $n \ll 0$.

§1. Associated primes of ideals and modules

Definition 1.1. Suppose M, N are R -submodules of some R -module M' . Then,

$$M :_R N := \{r \in R \mid rN \subseteq M\}.$$

Definition 1.2. Let M be an R -module and $x \in M$. If $\mathfrak{p} = 0 :_R x$ is a prime in R , then we say that \mathfrak{p} is an **associated prime** of M .

$$\text{Ass}_R(M) := \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} = 0 :_R x \text{ for some } x \in M\}.$$

Note that the x above would have to be nonzero since primes are proper ideals.

Definition 1.3. The elements of $\text{Ass}(M)$ which are not minimal in $\text{Ass}(M)$ are called **embedded primes**.

Definition 1.4. Fix $x \in M$. The map $\mu_x : R \rightarrow M$ defined by $r \mapsto rx$ is called the **homothety** by x .

Note that $\ker \mu_x = 0 :_R x = \text{ann}_R(x)$.

Proposition 1.5. A prime \mathfrak{p} is an associated prime of M iff R/\mathfrak{p} is isomorphic to a submodule of M .

Definition 1.6. $a \in R$ is a **zerodivisor** on M if $ax = 0$ for some $0 \neq x \in M$.

$$\mathcal{Z}(M) := \{a \in R \mid a \text{ is a zerodivisor on } M\}.$$

If a is not a zerodivisor, then a is called a **nonzerodivisor** on M or **M -regular**.

Note that a is a zerodivisor iff μ_a is not injective.

Proposition 1.7. Let R be Noetherian and $M \neq 0$ finitely generated R -module. Then,

1. the maximal elements among $\{(0 : x) \mid x \neq 0\}$ are prime. In particular, $\text{Ass } M \neq \emptyset$.
2. $\mathcal{Z}(M) = \bigcup_{\mathfrak{p} \in \text{Ass}(M)} \mathfrak{p}$.

Example 1.8. Let $R = k[x, y]$ for a field k and put $I = \langle x^2, xy \rangle$. Then, $\text{Ass}(R/I) = \{\langle x \rangle, \langle x, y \rangle\}$. Note that $\langle x \rangle$ is not maximal among the annihilators.

Proposition 1.9. Let $S \subseteq R$ be a multiplicatively closed set. Then,

1. $\text{Ass}_{S^{-1}R}(S^{-1}M) = \{S^{-1}\mathfrak{p} \mid \mathfrak{p} \in \text{Ass}(M), \mathfrak{p} \cap S = \emptyset\}$.
2. $\mathfrak{p} \in \text{Ass}_R(M) \Leftrightarrow \mathfrak{p}R_{\mathfrak{p}} \in \text{Ass}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}})$.

Definition 1.10. $\text{Supp}(M) := \{\mathfrak{p} \in \text{Spec}(R) \mid M_{\mathfrak{p}} \neq 0\}$.

Proposition 1.11. If M is finitely generated, then $\text{Supp}(M) = V(\text{ann}(M))$.

Proposition 1.12. If $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ is exact, then $\text{Supp } M = \text{Supp } N \cup \text{Supp } L$.

Proposition 1.13. Let L, K be f.g. R -modules. Then, $\text{Supp}(K \otimes_R L) = \text{Supp } L \cap \text{Supp } K$. In particular, $\text{Supp}(M/IM) = \text{Supp } M \cap V(I)$.

Proposition 1.14. $\text{Ass}(M) \subseteq \text{Supp}(M)$.

Note that if R is Noetherian and $I \trianglelefteq R$ is an ideal, then $\text{Ass}(R/I) \subseteq \text{Supp}(R/I) = V(I)$.

Assume that R and M are Noetherian from now.

Proposition 1.15. $\text{Ass } M$ and $\text{Supp } M$ have the same set of minimal primes.

Remark 1.16. Note that \mathfrak{p} is a minimal prime over \mathfrak{p}^n . That is, it is a minimal element of $V(\mathfrak{p}^n) = \text{Supp}(R/\mathfrak{p}^n)$ and hence, an element of $\text{Ass}(M/\mathfrak{p}^n)$.

Note that $V(\mathfrak{p}^n) = \text{Supp}(R/\mathfrak{p}^n)$ is true because of the Noetherian assumption.

Theorem 1.17. 1. There exists a sequence of R -submodules of M

$$(0) = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_{n-1} \subseteq M_n = M$$

such that $M_{i+1}/M_i \cong R/\mathfrak{p}_i$ for $\mathfrak{p}_i \in \text{Spec}(R)$.

2. Given any sequence as above, we have

$$\text{Ass } M \subseteq \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\} \subseteq \text{Supp } M.$$

In particular, $\text{Ass } M$ is always finite and hence, the set of minimal primes over any ideal is finite.

Definition 1.18. Let $N \leq M$ be a submodule such that $\text{Ass}(M/N) = \{\mathfrak{p}\}$. Then, M is called **\mathfrak{p} -primary**.

Definition 1.19. Let M be a module such that $\text{Ass } M = \{\mathfrak{p}\}$. Then, M is called **\mathfrak{p} -coprimary**.

Example 1.20. If $\mathfrak{m} \subseteq R$ is maximal, then \mathfrak{m}^n is \mathfrak{m} -primary for all $n \geq 1$. If $\mathfrak{p} \subseteq R$ is prime, then \mathfrak{p}^n need not be \mathfrak{p} -primary.

Proposition 1.21. If \mathfrak{q} is a \mathfrak{p} -primary ideal of R , then $\mathfrak{q}R_{\mathfrak{p}}$ is a $\mathfrak{p}R_{\mathfrak{p}}$ -primary ideal.

Proof. Note that $(R/\mathfrak{q})_{\mathfrak{p}} \cong R_{\mathfrak{p}}/\mathfrak{q}R_{\mathfrak{p}}$ as $R_{\mathfrak{p}}$ -modules. By Proposition 1.9, we see that

$$\begin{aligned} \mathfrak{a}R_{\mathfrak{p}} \in \text{Ass}_{R_{\mathfrak{p}}}(R_{\mathfrak{p}}/\mathfrak{q}R_{\mathfrak{p}}) &\Leftrightarrow \mathfrak{a}R_{\mathfrak{p}} \in \text{Ass}_{R_{\mathfrak{p}}}((R/\mathfrak{q})_{\mathfrak{p}}) \\ &\Leftrightarrow \mathfrak{a} \in \text{Ass}_R(R/\mathfrak{q}) = \{\mathfrak{p}\} \\ &\Leftrightarrow \mathfrak{a} = \mathfrak{p} \end{aligned}$$

and hence, $\mathfrak{q}R_{\mathfrak{p}}$ is $\mathfrak{p}R_{\mathfrak{p}}$ -primary. □

Definition 1.22. For $\mathfrak{a} \in R$, define $\mu_{\mathfrak{a}} : M \rightarrow M$ as $x \mapsto \mathfrak{a}x$.

Definition 1.23.

$$\begin{aligned} \text{nil}(M) &:= \{\mathfrak{a} \in R \mid \mu_{\mathfrak{a}} \text{ is nilpotent}\} \\ &= \{\mathfrak{a} \in R \mid \mathfrak{a}^n M = 0 \text{ for some } n\} \\ &= \sqrt{\text{ann}(M)} \end{aligned}$$

Proposition 1.24. If $\text{Ass}(M) = \{\mathfrak{p}\}$, then $\mathcal{Z}(M) = \text{nil}(M) = \sqrt{\text{ann}(M)}$.

Theorem 1.25. $|\text{Ass } M| = 1 \Leftrightarrow \mathcal{Z}(M) = \text{nil}(M)$.
If either condition holds, we have $\text{Ass } M = \{\sqrt{\text{ann}(M)}\}$.

Corollary 1.26. If $N \leq M$ is \mathfrak{p} -primary, then $\text{Ass}(M/N) = \{\sqrt{\text{ann}(M/N)}\}$.

Corollary 1.27. I is \mathfrak{p} -primary implies $\mathfrak{p} = \sqrt{I}$.

Remark 1.28. Note that if \sqrt{I} is prime, it does not imply that I is \sqrt{I} -primary.

Corollary 1.29. I is \mathfrak{p} -primary iff $\bigcup_{\mathfrak{p} \in \text{Ass}(R/I)} \mathfrak{p} = \mathcal{Z}(R/I) = \text{nil}(R/I) = I$.

Proposition 1.30. If N_1 and N_2 are \mathfrak{p} -primary, so is $N_1 \cap N_2$.

Definition 1.31. A submodule $N \leq M$ is called **reducible** if $N = N_1 \cap N_2$ with $N_1 \neq N \neq N_2$. It is called **irreducible** otherwise.

Proposition 1.32. Prime ideals are irreducible.

Theorem 1.33. Proper irreducible submodules are primary.

Theorem 1.34. Any proper submodule can be written as an intersection of finitely many irreducible submodules.

Corollary 1.35. Let R be a Noetherian ring and M a Noetherian R -module. If $N \subsetneq M$ is a proper R -submodule, then N can be written as

$$N = N_1 \cap \cdots \cap N_r,$$

where N_1, \dots, N_r are primary submodules.

The above is called a **primary decomposition** of N .

Definition 1.36. A primary decomposition is called **minimal** if $\text{Ass}(M/N_i) \neq \text{Ass}(M/N_j)$ for $i \neq j$.

It is called **irredundant** if N_i can be removed.

Theorem 1.37. If $N = N_1 \cap \cdots \cap N_r$ is an irredundant primary decomposition and $\text{Ass}(M/N_i) = \{p_i\}$, then $\text{Ass}(M/N) = \{p_1, \dots, p_r\}$.

Theorem 1.38. If p is a minimal associated prime of M/N , then the p -primary component of N is $\varphi_p^{-1}(Np)$, where $\varphi_p : M \rightarrow M_p$ is the natural map $x \mapsto \frac{x}{1}$.

In particular, the component corresponding to the minimal prime is uniquely determined.

§2. Artinian rings and Artinian modules

We now drop the assumption from the previous chapter of rings and modules being Noetherian.

Definition 2.1. An R -module M is called **Artinian** if every descending chain of R -submodules of M stabilises.

R is said to be an **Artinian ring** if R is Artinian as an R -module.

Proposition 2.2. Let k be a field and V a k -module, i.e., a k -vector space. Then, V is Artinian iff V is finite dimensional iff V is Noetherian.

Proposition 2.3. Let R be an Artinian ring.

1. If I is an ideal of R , then R/I is an Artinian ring.
2. If R is an integral domain, then R is a field.
3. More generally, every nonzerodivisor of R is a unit.
4. If $\mathfrak{p} \in \text{Spec}(R)$, then \mathfrak{p} is maximal. That is, $\text{Spec}(R) = \text{mSpec}(R)$.
Thus, $\mathcal{N}(R) = \mathcal{J}(R)$.
5. R has finitely many maximal ideals. (It may have infinitely many ideals, however.)
6. If $I \trianglelefteq R$, then $\text{Ass}(R/I) = \text{Supp}(R/I) = V(I)$.
7. If $N = \mathcal{N}(R)$, then there exists k such that $N^k = 0$.
8. Let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be an exact sequence. Then M is Artinian iff N and L are Artinian.
In particular, $\bigoplus_{i=1}^n M_i$ is Artinian iff each M_i is.
9. If M is a finitely generated R -module, then M is an Artinian R -module and $R/\text{ann}(M)$ is an Artinian ring.

Proposition 2.4. Let M be an R -module and $\mathfrak{m}_1, \dots, \mathfrak{m}_n \in \text{mSpec } R$ are maximal ideals such that $\mathfrak{m}_1 \cdots \mathfrak{m}_n M = 0$. Then,
 M is Noetherian $\Leftrightarrow M$ is Artinian.

Note that the maximal ideals above need not be distinct. Moreover, R is not assumed to be Artinian.

Proposition 2.5. Let R be an Artinian ring. Then, R is Noetherian ring.

Proposition 2.6. Let R be a Noetherian ring with $\text{Spec } R = \text{mSpec } R$. Then, R is an Artinian ring.

Proposition 2.7. If R is Artinian and M an Artinian R -module, then M is a Noetherian R -module. In particular, M is finitely generated.

Theorem 2.8. Let R be an Artinian ring. Then, there exist uniquely determined Artinian local rings R_1, \dots, R_n such that

$$R \cong R_1 \times \cdots \times R_n.$$

Definition 2.9. An R -module $M \neq 0$ is called **simple** if the only R -submodules of M are 0 and M .

Proposition 2.10. An R -module M is simple iff $M \cong R/\mathfrak{m}$ for some $\mathfrak{m} \in \text{mSpec } R$. The isomorphism is as R -modules. In particular, M is cyclic.

Lemma 2.11. A simple module is both Noetherian and Artinian.

Definition 2.12. Let M be an R -module. A series of the form

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{n-1} \subsetneq M_n = M$$

is called a **composition series** if M_{i+1}/M_i is simple for each i . n is called the **length** of this composition series.

Note that a composition series has finite length, by definition.

Theorem 2.13. M has a composition series $\Leftrightarrow M$ is Artinian and Noetherian.

Definition 2.14. Let $M \neq 0$ be an R -module. Define

$$l_R(M) := \min\{n \mid M \text{ has a composition series of length } n\}.$$

$l_R(M) = \infty$ if the set on the right is empty. $l_R(M)$ is called the **length** of M over R .

Note that if $R = k$ is a field, then the length of M is simply the dimension.

Definition 2.15. If $l_R(M) < \infty$, then M is called a **finite length** module.

Proposition 2.16. M is a finite length module iff M is Artinian and Noetherian.

Proposition 2.17. Let R be a Noetherian ring and M a finite length R -module. Then, $\text{Ass}(M) \subseteq \text{mSpec}(R)$.

Proposition 2.18. Let M be a finite length module and $N \leq M$. Then, N also has finite length and $l_R(N) \leq l_R(M)$ with equality iff $N = M$.

Theorem 2.19 (Jordan-Hölder). Every composition series of a finite length module M has the same length.

Now, if

$$\begin{aligned} 0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{n-1} \subsetneq M_n = M, \\ 0 = N_0 \subsetneq N_1 \subsetneq \cdots \subsetneq N_{n-1} \subsetneq N_n = M \end{aligned}$$

are two composition series of M , then there exists a permutation $\sigma \in S_n$ such that

$$M_i/M_{i-1} \cong N_{\pi(i)}/N_{\pi(i)-1}$$

for all $1 \leq i \leq n$. In other words, the quotients that appear are the same.

Proposition 2.20. Let M be a finite length module. Any series of the form

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{n-1} \subsetneq M_n = M$$

is actually a composition series.

Proposition 2.21. Let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be an exact sequence. Then, $l_R(M) = l_R(N) + l_R(L)$.

Note that M is finite length iff N and L both are.

Proposition 2.22. If R is Noetherian and M a finite length R -module, then $\text{Ass}(M) \subseteq \text{mSpec}(R)$.

§3. Integral Extensions of Rings

Definition 3.1. Let $R \subseteq S$ be nonzero commutative rings. An element $s \in S$ is called **integral** over R if there exists a monic polynomial $f(x) \in R[x]$ such that $f(s) = 0$.

Let

$$T = \{s \in S \mid s \text{ is integral over } R\}.$$

T is called the **integral closure** of R in S .

If R is an integral domain and $S = Q(R)$, then T is called the **normalisation** of R . R is called **normal** or **integrally closed** if $T = R$.

Recall that if R is an integral domain, then $Q(R)$ denotes the field of fractions of R .

We shall shortly show that T is a subring of S .

Theorem 3.2. If R is a UFD, then R is integrally closed. In other words, UFDs are normal.

The converse is not true.

Theorem 3.3 (Cayley-Hamilton). Let $I \trianglelefteq R$ be an ideal and M a finitely generated R -module. Let $\varphi : M \rightarrow M$ be an R -endomorphism such that $\varphi(M) \subseteq IM$. Then, φ satisfies a monic polynomial of the form

$$x^n + a_1x^{n-1} + \cdots + a_n$$

with $a_1, \dots, a_n \in I$.

Corollary 3.4 (Nakayama). Suppose M is finitely generated over R and $I \trianglelefteq R$ is such that $M = IM$. Then, there exists $a \in I$ such that $(1 + a)M = 0$. In particular, if $I \subseteq \mathcal{J}(R)$, then $M = 0$.

Corollary 3.5. If $\varphi : M \rightarrow M$ is a surjective R -linear map, then φ is an isomorphism. (M is finitely generated as usual.)

Corollary 3.6. Suppose $M \cong R^n$. Then, any set of n generators is linearly independent.

Corollary 3.7. Let R be a nonzero commutative ring. Then, $R^n \cong R^m$ (as R -modules) iff

$n = m$.

Theorem 3.8. Let $R \subseteq S$ be nonzero commutative rings and $s \in S$. TFAE:

1. s is integral over R .
2. $R[s]$ is a finitely generated R -module.
3. There exists a subring T such that $R[s] \subseteq T \subseteq S$ and T is a finitely generated R -module.

Theorem 3.9. Let $R \subseteq S$ be a ring extension and $T = \overline{R}^S$ the integral closure of R in S . Then, T is a subring of S .

Proposition 3.10. If $R \subseteq T$ and $T \subseteq S$ are integral extensions, then so is $R \subseteq S$.

Corollary 3.11. If T is the integral closure of R in S , then the integral closure of T in S is T .

Symbolically, if $T = \overline{R}^S$, then $\overline{T}^S = T$.

Note that if $R \subseteq S$ is a ring extension and $I \trianglelefteq S$ is an ideal, then $I^c = R \cap I$ is an ideal of R . (Called the contraction.) Also, one has the natural inclusion and projection maps as

$$R \xrightarrow{i} S \xrightarrow{\pi} S/I.$$

Then, $I^c = \ker(\pi \circ i)$ and hence, R/I^c is isomorphic to a subring of S/I . We denote this inclusion by writing $R/I^c \hookrightarrow S/I$.

Proposition 3.12. If $R \subseteq S$ is an integral extension, then so is $R/I^c \hookrightarrow S/I$.

Definition 3.13. Suppose $\varphi : R \rightarrow S$ is a ring map. Then, φ is called **integral** if $\varphi(R) \subseteq S$ is an integral extension.

Proposition 3.14. Let $U \subseteq R$ be a multiplicatively closed subset and let $R \subseteq S$ be an integral extension. Then, $U^{-1}R \subseteq U^{-1}S$ is an integral extension.

Proposition 3.15. Let R be an integral domain. TFAE:

1. R is integrally closed (normal).
2. R_p is integrally closed for all $p \in \text{Spec}(R)$.
3. R_m is integrally closed for all $m \in \text{mSpec}(R)$.

Lemma 3.16. Let $R \subseteq S$ be an integral extension of integral domains. Then, R is a field $\Leftrightarrow S$ is a field.

Corollary 3.17. Let $R \subseteq S$ be rings (not necessarily domains) and $q \in \text{Spec } S$. Define $p := R \cap q$. Then, $p \in \text{mSpec } R \Leftrightarrow q \in \text{mSpec } S$.


In particular, given an integral extension, the contraction of a maximal ideal is maximal.

Definition 3.18. Let $R \subseteq S$ be rings. Suppose $Q \in \text{Spec } S$ and $P \in \text{Spec } R$. Q is said to **lie over** P if $Q^c = Q \cap R = P$.


Theorem 3.19 (Lying over theorem). Let $R \subseteq S$ be an integral extension of rings and $p \in \text{Spec } R$. Then, there exists $q \in \text{Spec } S$ such that $q \cap R = p$.

In other words: Given an integral extension, there is always a prime lying over a prime.


Theorem 3.20 (Going up theorem). Let $R \subseteq S$ be an integral extension. Let $p_1, p_2 \in \text{Spec } R$ with $p_1 \subseteq p_2$ and $q_1 \in \text{Spec } S$ be such that $q_1 \cap R = p_1$. Then, there exists $q_2 \in \text{Spec } S$ such that $q_1 \subseteq q_2$ and $q_2 \cap R = p_2$.



In fact, inductively, we see that any chain above can be “completed.”



Proposition 3.21 (Incompatibility (INC)). Let $R \subseteq S$ be an integral extension of rings. Let $Q_1, Q_2 \in \text{Spec } S$ lie over $P \in \text{Spec } R$. If Q_1 and Q_2 are distinct, then they are incomparable. That is, $Q_1 \neq Q_2 \implies Q_1 \not\subseteq Q_2$ and $Q_2 \not\subseteq Q_1$.




Lemma 3.22. Let $f : R \rightarrow S$ be any ring homomorphism and $P \in \text{Spec } R$. TFAE:

1. $P^{ec} = f^{-1}(f(P)S) = P$, and
2. $\exists Q \in \text{Spec } S$ such that $Q^c = P$. That is, a prime lies over P .


Note that the above is a general fact, no assumptions of integral extensions.

Theorem 3.23 (Going down theorem). Let R be a normal domain, S an integral domain and $R \subseteq S$ be an integral extension.

Given $P_0, P_1 \in \text{Spec } R$ with $P_0 \supseteq P_1$ and a prime $Q_0 \in \text{Spec } S$ lying over P_0 , there exists a prime $Q_1 \in \text{Spec } S$ lying over P_1 with $Q_0 \supseteq Q_1$.



In fact, inductively, we see that any chain above can be “completed.”



Theorem 3.24. Let R be a Noetherian normal domain with quotient field K . Let $K \subseteq L$ be a separable extension. Then, \bar{R}^L is a finite R -module. In particular, it is a Noetherian ring.

§4. Dimension Theory of Affine Algebra over Fields

Lemma 4.1 (Artin-Tate Lemma). Let $R \subseteq S \subseteq T$ be rings. Suppose

1. R is Noetherian,
2. T is a finitely generated S module,
3. T is a finitely generated R algebra.

$$\begin{array}{c}
 R[t_1, \dots, t_s] = T = St'_1 + \dots + St'_k \\
 | \\
 S \\
 | \\
 R
 \end{array}$$

Then, S is a finitely generated R -algebra. In other words, there exist $s_1, \dots, s_n \in S$ such that $S = R[s_1, \dots, s_n]$.

In particular, S is Noetherian.

Definition 4.2. Let k be a field. An **affine k -algebra** is a ring of the form $R = k[x_1, \dots, x_n]/I$ for some ideal $I \subseteq k[x_1, \dots, x_n]$.

Lemma 4.3 (Zariski). Let k be any field and $R = k[x_1, \dots, x_n]/I$ be an affine k -algebra which is also a field. (That is, I is maximal.)

Then, R is an algebraic extension of k .

Corollary 4.4. Let $\varphi : R \rightarrow S$ be a ring homomorphism, where R and S are affine k -algebras. Let $\mathfrak{m} \in \text{mSpec}(S)$. Then, $\varphi^{-1}(\mathfrak{m}) \in \text{mSpec}(R)$.

(We had used the fact that if we have an algebraic extension $K \subseteq F$ of fields and an integral domain R such that $K \subseteq R \subseteq F$, then R is a field.)

Theorem 4.5 (Weak Nullstellensatz). If k is algebraically closed, then maximal ideals $\mathfrak{m} \in \text{mSpec } k[x_1, \dots, x_n]$ are precisely those of the form $\mathfrak{m}_a = (x_1 - a_1, \dots, x_n - a_n)$ for some $(a_1, \dots, a_n) \in k^n$.

Corollary 4.6 (Criterion for solvability). Let $p_1(x_1, \dots, x_n), \dots, p_s(x_1, \dots, x_n)$ be polyno-

mials in $k[x_1, \dots, x_n]$. Then, the polynomials have a common solution iff the ideal generated by them is not the whole ring.

Remark 4.7. In fact, one need not assume $s < \infty$ in the above.

Definition 4.8. Given a field k , \mathbb{A}_k^n denotes the **affine n -space over k** . It is simply the set k^n without any vector space structure.

Given any ideal $I \trianglelefteq k[x_1, \dots, x_n]$, we define the **zero set of I** as

$$\mathcal{Z}(I) = \{\underline{a} \in \mathbb{A}_k^n : f(\underline{a}) = 0 \text{ for all } f \in I\} \subseteq \mathbb{A}_k^n.$$

A subset of \mathbb{A}_k^n which is the zero set of some ideal is called an **algebraic set**.

Given an algebraic set $X \subseteq \mathbb{A}_k^n$, we define the **ideal of X** as

$$\mathcal{I}(X) = \{f \in k[x_1, \dots, x_n] : f(x) = 0 \text{ for all } x \in X\} \subseteq k[x_1, \dots, x_n].$$

Remark 4.9. An ideal of an algebraic set is always a radical ideal.

Theorem 4.10 (Strong Nullstellensatz). If k is algebraically closed and $I \trianglelefteq k[x_1, \dots, x_n] = S$ an ideal, then $\mathcal{I}(\mathcal{Z}(I)) = \sqrt{I}$.

In particular, there is a bijection

$$\{\text{radical ideals in } S\} \leftrightarrow \{\text{algebraic subsets in } \mathbb{A}_k^n\}.$$

Definition 4.11. Given a polynomial $f \in k[x_1, \dots, x_n]$, we can write

$$f = \sum_{\alpha \in (\mathbb{N} \cup \{0\})^n} \alpha_\alpha x^\alpha.$$

If $\alpha_\alpha \neq 0$, we say that x^α is a **term** of f .

Writing $\alpha = (\alpha_1, \dots, \alpha_n)$, $|\alpha|$ denotes the maximum of $\alpha_1, \dots, \alpha_n$.

Proposition 4.12. Let k be any field. Let $f \in S = k[x_1, \dots, x_n]$ be a non-constant polyno-

mial. Let

$$N > \max\{|\alpha| : \alpha \in (\mathbb{N} \cup \{0\})^n, x^\alpha \text{ is a term of } f\}.$$

Without loss of generality, we may assume that x_n appears non-trivially in some term of f . Define the map $\Phi : S \rightarrow S$ by identity on k and

$$x_i \mapsto \begin{cases} x_i - x_n^{N^i} & i \neq n, \\ x_n & i = n. \end{cases}$$

Then, Φ is an automorphism such that $\Phi(f)$ is monic in x_n , up to a constant. That is,

$$\Phi(f) = cx_n^r + g_1x_n^{r-1} + \cdots + g_n,$$

where $0 \neq c \in k$ and $g_1, \dots, g_n \in k[x_1, \dots, x_{n-1}]$.

Theorem 4.13 (Noetherian Normalisation Lemma). Let $R = k[\theta_1, \dots, \theta_n]$ be an affine k -algebra. Then, there exist algebraically independent elements $z_1, \dots, z_d \in R$ such that $k[z_1, \dots, z_d] \subseteq R$ is an integral extension.

$$\begin{array}{c} R \\ \left| \text{integral} \right. \\ k[z_1, \dots, z_d] = S. \end{array}$$

In particular, R is a finite S -module.

Corollary 4.14. Let R be an affine k -algebra and $I \trianglelefteq R$ an ideal. Then

$$\sqrt{I} = \bigcap_{\mathfrak{m}: I \subseteq \mathfrak{m} \in \text{Spec}(R)} \mathfrak{m}.$$

In particular, $\mathcal{N}(R) = \mathcal{J}(R)$.

Definition 4.15. A **saturated chain of prime ideals** is a chain

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

of prime ideals such that no prime ideal can be inserted strictly in between anywhere above. (In other words, there exists no $i \in \{0, \dots, n-1\}$ and no $\mathfrak{q} \in \text{Spec}(R)$ such that $\mathfrak{p}_i \subsetneq \mathfrak{q} \subsetneq \mathfrak{p}_{i+1}$.)

The **length** of the above chain is n . The **Krull dimension** of R is defined as

$$\dim(R) = \sup\{n : \exists \text{ a saturated chain of length } n\}.$$

$\dim(R)$ may be ∞ even if R is Noetherian.

Definition 4.16. Given a prime ideal $\mathfrak{p} \subseteq R$, the **height** of \mathfrak{p} is defined as

$$\text{ht}(\mathfrak{p}) = \dim(R_{\mathfrak{p}}).$$

Example 4.17. Here are some examples.

1. If R is Artinian, then $\dim(R) = 0$. In particular, $\dim(k) = 0$.
2. $\dim(\mathbb{Z}) = 1$.
3. $\dim(k[X]) = 1$.
4. In general, if R is a PID and not a field, then $\dim(R) = 1$.

Proposition 4.18. Let $R \subseteq S$ be an integral extension of rings. Then,

1. $\dim(R) = \dim(S)$.
2. If $I \triangleleft S$ is a proper ideal, then $\dim(S/I) = \dim(R/I \cap R)$.
3. Suppose S is integral and R normal. Let $Q \in \text{Spec}(S)$. Then, $\dim(S_Q) = \dim(R_{Q \cap R})$.

Theorem 4.19. Let R be an affine algebra over a field k . Let $z_1, \dots, z_d \in R$ be such that $S = k[z_1, \dots, z_d] \subseteq R$ is an integral extension. (Exists by NNL.)

Then,

1. $\dim(R) = d = \dim(k[z_1, \dots, z_d])$.
2. Any maximal saturated chain of prime ideals in R has length d .

Remark 4.20. The above shows that the d in **Noetherian Normalisation Lemma** is determined uniquely. Moreover, it shows that the dimension of polynomial ring in d variables over a field is d .

§5. Graded Rings and Graded Modules

Definition 5.1. Let R be any commutative ring. (In particular, it is an additive subgroup.) Let $\{R_n : n = 0, 1, \dots\}$ be a sequence of additive subgroups with the properties that

$$R = \bigoplus_{n \geq 0} R_n \quad \text{and} \quad R_n R_m \subseteq R_{n+m} \text{ for all } n, m \geq 0.$$

Then, R is called a **graded ring** with **grading** $(R_n)_n$. Any $x \in R_n$ is called **homogeneous of degree n** .

Remark 5.2. Note that for $n = m = 0$, we have $R_0 R_0 \subseteq R_0$. Thus, R_0 is closed under multiplication as well. Moreover, writing $1 = r_0 + \dots + r_n$ for $r_i \in R_i$ and noting that $1^2 = 1$ gives that $n = 0$ and $1 \in R_0$.

Thus, R_0 is actually a subring of R . In particular, R is an R_0 -module. Moreover, $R_0 R_n \subseteq R_n$ gives us that each R_n is an R_0 -module.

Example 5.3. Consider $R = k[X_1, \dots, X_r]$ and let R_n be the k vector space generated by monomials of degree n . Then, R is a graded ring with grading $(R_n)_n$.

Definition 5.4. Let R be a graded ring with grading $(R_n)_{n \geq 0}$. An R -module M is called **graded** if there exists a sequence of additive subgroups $(M_n)_{n \geq 0}$ such that

$$M = \bigoplus_{n \in \mathbb{Z}} M_n \quad \text{and} \quad R_m M_n \subseteq M_{n+m} \text{ for all } m, n \geq 0.$$

Remark 5.5. Note that M_n are not necessarily R -submodules of M . As an example, if $r \in R_1$ and $x \in M_n$, then $rx \in M_{n+1}$. If $rx \neq 0$, then $rx \notin M_n$ and so, M_n is not closed under scalar multiplication.

However, one notes that each M_n is an R_0 -submodule of M . (Note that since R_0 is a subring of R , we may regard M as an R_0 -module as well.)

Note that in the above we have used \mathbb{N}_0 for grading. However, we may use \mathbb{N} or \mathbb{Z} as well. In fact, one may use any monoid or even semigroup.

Definition 5.6. Let R be a graded ring and M a graded R -module. A submodule $N \leq M$

is called a **graded submodule** if N is generated by homogeneous elements of M .

Theorem 5.7 (Characterisation of graded submodules). Let $R = \bigoplus_{n \geq 0} R_n$ be a graded ring and $M = \bigoplus_{n \geq 0} M_n$ a graded R -module and $N \leq M$ a submodule. TFAE:

1. N is a graded R -submodule of M , that is, N is generated by homogeneous elements.
2. $N = \bigoplus_{n \geq 0} (N \cap M_n)$.
3. If $y \in N$ and $y = y_0 + \cdots + y_n$, where $y_i \in M_i$, then $y_i \in N$.

The second point says that N can be considered a graded R -submodule by itself. The third says that if we write an element of N as a sum of homogeneous elements of different degrees, then each such element must be in N itself.

Example 5.8. Consider $R = M = k[x]$, with the usual grading as in the previous example. Then, the submodule $I = \langle x \rangle$ is a graded R -submodule of M since it is generated by a homogeneous element.

On the other hand, the submodule $J = \langle x - 1 \rangle$ is *not* graded because we have $x - 1 \in J$ and $x, 1$ are homogeneous of different degrees but $1 \notin J$.

Theorem 5.9 (Characterisation of Noetherian graded rings). Let $R = \bigoplus_{n \geq 0} R_n$ be a graded ring. TFAE:

1. R is Noetherian.
2. R_0 is Noetherian and R is a finitely generated R_0 -algebra, i.e., $R = R_0[r_1, \dots, r_n]$.

Definition 5.10. The sum $R_+ = \bigoplus_{i \geq 1} R_i$ is an ideal of R , called the irrelevant ideal.

Definition 5.11. Let R be a commutative ring and $F = \{I_n\}_{n \geq 0}$ a filtration of ideals as

$$R = I_0 \supseteq I_1 \supseteq I_2 \supseteq \cdots$$

satisfying $I_n I_m \subseteq I_{n+m}$.

Let t be an indeterminate. We define the **Rees ring of F** as

$$\mathcal{R}(F) = \bigoplus_{n \geq 0} I_n t^n \subseteq R[t].$$

$\mathcal{R}(F)$ is a graded ring with grading $(I_n t^n)_{n \geq 0}$.

The fact that it is graded follows from the condition that $I_n I_m \subseteq I_{n+m}$.
 A special case of the above is when we take $I_n = I^n$ for some fixed ideal $I \trianglelefteq R$.

Definition 5.12. The **Rees ring of an ideal** I is defined as

$$\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n t^n = \left\{ \sum a_i t^i : a_i \in I^i \right\}.$$

(Convention: $I^0 = R$.)

In the following, we have the following notation: Let R be a ring, $I \trianglelefteq R$ and ideal and M and R -module.

Definition 5.13. Let $M = M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots$ be a filtration of submodules. Then this filtration is called an **I-filtration** if

$$IM_n \subseteq M_{n+1} \text{ for all } n \in \mathbb{N}.$$

$\bigoplus_{n \geq 0} M_n t^n \subseteq M[t]$ is a graded $\mathcal{R}(I)$ -module since $I^n M_m \subseteq M_{n+m}$ for all $n, m \geq 0$.

The filtration is called **I-stable** if

$$IM_n = M_{n+1} \text{ for } n \gg 0.$$

That is, there exists $N \in \mathbb{N}$ such that $IM_n = M_{n+1}$ for all $n > N$.

Example 5.14. $\{I^n M\}_{n \geq 0}$ is an I-stable filtration.

Definition 5.15. Given filtrations $F = \{I_n\}_{n \geq 0}$ of ideals of R and $G = \{M_n\}_{n \geq 0}$ of R -submodules of M , we say that G is **F-compatible** if $I_n M_m \subseteq M_{n+m}$ for all $n, m \geq 0$.

Example 5.16. $\{I^n M\}_{n \geq 0}$ is $\{I^n\}_{n \geq 0}$ -compatible.

Theorem 5.17. Suppose R is Noetherian, $I \trianglelefteq R$ an ideal, M a finitely generated R -module and G a filtration of R -submodules of M which is $\{I^n\}_{n \geq 0}$ -compatible. Then,

$$\mathcal{R}(G) \text{ is a finitely generated } \mathcal{R}(I)\text{-module} \Leftrightarrow G \text{ is I-stable.}$$

The next two theorems (ARL and KIT) were obtained as corollaries.

Theorem 5.18 (Generalised Artin-Rees Lemma). Suppose R is Noetherian, $I \trianglelefteq R$ an ideal, M a finitely generated R -module, $N \leq M$ an R -submodule, and $\{M_n\}_{n \geq 0}$ is an I -stable filtration. Then, $\{N \cap M_n\}_{n \geq 0}$ is I -stable.

In other words, if $IM_n = M_{n+1}$ for $n \gg 0$, then $I(M_n \cap N) = M_{n+1} \cap N$ for $n \gg 0$.

Corollary 5.19 (Artin-Rees Lemma). Suppose R is Noetherian, $I \trianglelefteq R$ an ideal, M a finitely generated R -module, $N \leq M$ an R -submodule. Then,

$$N \cap I^n M = I^{n-k}(N \cap I^k M) \text{ for } n \gg 0.$$

Theorem 5.20 (Krull Intersection Theorem). Let R be a Noetherian ring, $I \trianglelefteq R$ an ideal, and M a finitely generated R -module. Define the submodule $N \leq M$ as

$$N := \bigcap_{n \geq 1} I^n M.$$

Then,

1. There exists $a \in I$ such that $(1 + a)N = 0$.
2. If $I \subseteq \mathcal{J}(R)$, then $N = \bigcap_{n \geq 1} I^n M = 0$.
3. If R is a Noetherian domain and $I \neq R$, then $\bigcap_{n \geq 1} I^n = 0$.
4. If R is local and $I \neq R$, then $N = \bigcap_{n \geq 1} I^n M = 0$.

Example 5.21. KIT need not be true if R is not Noetherian. Consider the ring $R = \mathcal{C}^\infty(\mathbb{R})$, the ring of smooth real valued functions defined on \mathbb{R} with pointwise operations. This ring is not Noetherian. We construct a maximal ideal such that the intersection of its powers is not zero.

Consider the map $\varphi : R \rightarrow \mathbb{R}$ given by $f \mapsto f(0)$. This is a surjective ring homomorphism. Since \mathbb{R} is a field, we see that $\mathfrak{m} := \ker \varphi$ is maximal.

Let $x \in R$ denote the identity function.

Claim 1. $\mathfrak{m} = (x)$.

Proof. \supseteq is clear since $x(0) = 0$.

(\subseteq) Let $f \in \mathfrak{m}$. Consider the function $g : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$g(t) := \begin{cases} f(t)/t & t \neq 0, \\ f'(0) & t = 0. \end{cases}$$

Then, one sees that g is smooth. Clearly, g is infinitely differentiable on $\mathbb{R} \setminus \{0\}$. For 0 , one may use L'Hôpital's rule to successively compute the derivatives (and show their existence).

Clearly, $xg = f$ and hence, $f \in (x)$, as desired. \square

In particular, we have $\mathfrak{m}^n = (x^n)$. Note that x^n is the function $t \mapsto t^n$. We now identify this ideal.

Claim 2. $(x^n) = \{f \in \mathbb{R} : f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0\}$.

Proof. As before \subseteq is clear since $(x^n)' = nx^{n-1}$.

(\supseteq) The idea is as before. Let $g \in \mathbb{R}$ be such that $f(0) = \dots = f^{(n-1)}(0) = 0$. Define $g : \mathbb{R} \rightarrow \mathbb{R}$ by

$$g(t) := \begin{cases} f(t)/t^n & t \neq 0, \\ f^{(n)}(0)/n! & t = 0. \end{cases}$$

As earlier, we get $g \in \mathbb{R}$ and $f = x^n g \in (x^n)$. \square

Claim 3. $\bigcap_{n \geq 1} \mathfrak{m}^n \neq 0$.

Proof. Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) := \begin{cases} e^{-1/x^2} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Then, $f \in \mathbb{R}$ and $f^{(n)}(0) = 0$ for all n . However, f is not the zero function. Thus, $0 \neq f \in \bigcap_{n \geq 1} \mathfrak{m}^n$, as desired. \square

Definition 5.22. Given an ideal $I \trianglelefteq R$, we define the ideal I^* as

$$I^* = \bigoplus_{n \geq 0} (I \cap R_n).$$

Note that $I^* \subseteq I$.

Recall that $I = I^* \Leftrightarrow I$ is generated by homogeneous elements. (**Characterisation of graded submodules.**)

Proposition 5.23. Let $R = \bigoplus_{n \geq 0} R_n$ be a graded ring.

1. If $\mathfrak{p} \in \text{Spec}(R)$, then $\mathfrak{p}^* \in \text{Spec}(R)$.
2. Any minimal prime of R is a graded ideal.
3. Maximal graded ideals of R are precisely of the form $\mathfrak{m} \oplus R_1 \oplus R_2 \oplus \cdots$ for $\mathfrak{m} \in \text{mSpec}(R)$.
4. If R is Noetherian and $M = \bigoplus_{n \geq 0} M_n$ graded, then $\mathfrak{p} \in \text{Ass}_R(M)$ is graded and $\mathfrak{p} = \text{ann}_R(x)$ for some *homogeneous* element $x \in M$.
5. If the nonzero *homogeneous* elements of R are nonzerodivisors, then R is an integral domain.
6. Let $\mathfrak{p} \neq R$ be a graded ideal. If for all *homogeneous* elements $a, b \in R$ the following holds:

$$ab \in \mathfrak{p} \implies a \in \mathfrak{p} \text{ or } b \in \mathfrak{p},$$

then \mathfrak{p} is a prime ideal.

§6. Dimension Theory of Finite Modules over Noetherian Local Rings

Recall that by “finite module,” we mean “finitely generated modules.”

In the following, R will denote a Noetherian ring and M a finitely generated R -module.

Definition 6.1. The **Krull dimension** of $M \neq 0$ is defined as $\dim(M) := \dim(R/\text{ann}(M))$. If $M = 0$, we define $\dim(M) = -1$.

(Recall that we had already defined the Krull dimension of a ring in Definition 4.15.)

Caution: Note that if R is a field and V a finite dimensional vector space over R , then $\dim(V)$ will **not** be the usual vector space dimension. If $V \neq 0$, then (Krull) $\dim(V) = 0$ since $R/\text{ann}(M) = R$ and $\dim(\text{field}) = 0$. On the other hand, if $V = 0$, then (Krull) $\dim(V) = -1$, by the above convention.

Definition 6.2. The **Chevalley dimension** of $M \neq 0$ is defined as

$$c(M) := \inf\{n : \exists x_1, \dots, x_n \text{ s.t. } l_R(M/(x_1, \dots, x_n)M) < \infty\}.$$

For $M = 0$, we define $c(M) := -1$.

Note that if $M \neq 0$ and M has finite length, then $c(M) = 0$.

Proposition 6.3. Let $R = \bigoplus_{n \geq 0} R_n$ be a graded ring Noetherian and $M = \bigoplus_{n \in \mathbb{Z}} M_n$ a finitely generated graded R -module. Then,

1. $M_n = 0$ for $n \ll 0$,
2. each M_n is a finitely generated R_0 -submodule.

In particular, if R_0 is Artinian as well, then $l_{R_0}(M_n) < \infty$ for all $n \in \mathbb{Z}$.

Definition 6.4. Given any ring R , we define the **Laurent series ring over R in the variable x** as the localisation of $R[[x]]$ at the set $\{1, x, x^2, \dots\}$.

In other words, it is the set of (formal) series of the form

$$\sum_{n=N}^{\infty} a_n x^n$$

for some $N \in \mathbb{Z}$ and some $a_n \in R$.

We denote this ring by $R[x, x^{-1}]$.

$R[x, x^{-1}]$ is the subring of $R[[x, x^{-1}]]$ which has only finitely many $a_n \neq 0$.

Definition 6.5. A function $f : \mathbb{Z} \rightarrow \mathbb{Q}$ is called **polynomial-like** if there exists $g(x) \in \mathbb{Q}[x]$ such that $f(n) = g(n)$ for $n \gg 0$. In this case, we define $\deg(f) = \deg(g(x))$ and call g the **corresponding polynomial** of f .

Remark 6.6. Note that if f is polynomial-like, then the polynomial g in the definition above is uniquely determined. Indeed, if h is another such polynomial, then we have $(g - h)(n) = 0$ for n sufficiently large. Thus, $g = h$.

This justifies the degree and corresponding polynomial being well-defined.

Definition 6.7. Let R be a Noetherian graded ring such that R_0 is Artinian. Let M be a finite graded R -module. We define the **Hilbert series** of M as

$$H(M, \lambda) = \sum_{n \in \mathbb{Z}} l_{R_0}(M_n) \lambda^n \in \mathbb{Z}[[\lambda, \lambda^{-1}]].$$

Theorem 6.8 (Hilbert-Serre). Let R be a Noetherian graded ring such that R_0 is Artinian. Let M be a finite graded R -module.

Suppose $x_1, \dots, x_r \in R$ are such that $R = R_0[x_1, \dots, x_r]$ with $d_i := \deg(x_i) \geq 1$. Then,

$$H(M, \lambda) = \frac{f(\lambda, \lambda^{-1})}{\prod_{i=1}^r (1 - \lambda^{d_i})}$$

for some $f(\lambda, \lambda^{-1}) \in \mathbb{Z}[[\lambda, \lambda^{-1}]]$.

Corollary 6.9. In the same setup as earlier, $n \mapsto l_{R_0}(M_n)$ is a polynomial like function.

Definition 6.10. Let R be a Noetherian graded ring such that R_0 is Artinian. Let M be a finite graded R -module.

The $n \mapsto l_{R_0}(M_n)$ is (loosely) called the **Hilbert polynomial** of M .

Note that the above is a polynomial-like function and not necessarily given by a poly-

mial itself.