

# Extending Riemann maps to the boundary

Aryaman Maithani

IIT Bombay

16th March 2022

# Notations and Conventions

- 1  $\mathbb{D}$  will denote the open unit disc.  $S^1 := \partial\mathbb{D}$  is the unit circle.
- 2  $\mathcal{O}^\infty$  denotes the set of bounded holomorphic functions on  $\mathbb{D}$ .
- 3  $\Omega$  will always denote a nonempty, open, bounded, and simply-connected subset of  $\mathbb{C}$ .
- 4 Recall that a Riemann mapping of  $\Omega$  onto  $\mathbb{D}$  is simply a biholomorphism  $\Omega \rightarrow \mathbb{D}$ .
- 5 A curve shall mean a continuous function with domain  $[0, 1]$ . Typically,  $\gamma$  will be a curve such that  $\gamma([0, 1)) \subseteq \Omega$  and  $\gamma(1) \in \partial\Omega$ . Similarly,  $\Gamma$  will be a curve such that  $\Gamma([0, 1)) \subseteq \mathbb{D}$  and  $\Gamma(1) \in \partial\mathbb{D}$ .

Let  $\Omega$  be a bounded simply-connected domain in  $\mathbb{C}$ . By the Riemann Mapping Theorem, we know that there exists a biholomorphism  $f : \Omega \rightarrow \mathbb{D}$ . The following is a natural question.

## Question

Can  $f$  be continuously extended up to  $\overline{\Omega}$ ?

The obvious way to extend  $f$  is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique and we must have  $f(\overline{\Omega}) \subseteq \overline{\mathbb{D}}$ . This must force  $f(\overline{\Omega}) = \overline{\mathbb{D}}$ . (Why?)

Thus, it also makes sense to ask whether *the* extension is a homeomorphism onto  $\overline{\mathbb{D}}$ .

What we shall see is the following: By imposing a *simple* topological restriction on  $\Omega$ , one gets that *any* biholomorphism  $\Omega \rightarrow \mathbb{D}$  can be extended to a continuous *injective* map  $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$ . Moreover, this will be a *homeomorphism*.

## Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended  $f$  to  $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$ , we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

As  $\tilde{f}(\overline{\Omega})$  is compact, we have  $\tilde{f}(\overline{\Omega}) = \overline{\mathbb{D}}$ .

Furthermore, if  $\tilde{f}$  is an injection, then compactness again tells us that  $\tilde{f}$  is a homeomorphism (as  $\tilde{f}$  is a bijection).

## Definition 2

A boundary point  $\beta$  of  $\Omega$  is called a **simple boundary point** if  $\beta$  has the following property:

For every sequence  $(\alpha_n)_n$  in  $\Omega$  such that  $\alpha_n \rightarrow \beta$  as  $n \rightarrow \infty$ , there exists a curve  $\gamma$  and a strictly increasing sequence  $(t_n)_n$  in  $(0, 1)$  such that


$$t_n \rightarrow 1, \gamma(t_n) = \alpha_n \ (n = 1, 2, \dots), \gamma([0, 1)) \in \Omega.$$

$\gamma(1) = \beta$  follows by continuity.

In words: there is a curve in  $\Omega$  which passes through  $\alpha_n$  and ends at  $\beta$ .

# (Non-)Examples

- 1 Every boundary point of  $\mathbb{D}$  is a simple boundary point.
- 2 Let  $\Omega := \mathbb{D} \setminus [0, 1)$ .



The nonzero boundary points of  $\Omega$  lying on the real axis are not simple. Note that  $\Omega$  is indeed bounded and simply-connected and thus, biholomorphic to  $\mathbb{D}$ . However,  $\partial\Omega$  is clearly not homeomorphic to  $\partial\mathbb{D}$  and thus, no biholomorphism can be extended to a homeomorphism  $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$ .

## Theorem 3 (Helper Theorem)

Let  $\Omega$  be a bounded simply-connected domain, and let  $f$  be a Riemann mapping of  $\Omega$  onto  $\mathbb{D}$ .

- 1 If  $\beta$  is a simple boundary point of  $\Omega$ , then  $f$  has a continuous extension to  $\Omega \cup \{\beta\}$ . If  $f$  is so extended, then  $|f(\beta)| = 1$ .
- 2 If  $\beta_1$  and  $\beta_2$  are distinct simple boundary points of  $\Omega$  and if  $f$  is continuously extended to  $\Omega \cup \{\beta_1, \beta_2\}$ , then  $f(\beta_1) \neq f(\beta_2)$ .

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in 1 is unique and would have to be attained as follows: given a sequence  $(\alpha_n)_n$  in  $\Omega$  converging to  $\beta$ , we have  $f(\beta) := \lim f(\alpha_n)$ . Once we show that this limit (exists and) is independent of the sequence  $(\alpha_n)$ , we would have shown continuity.

# The Main Extension Theorem

## Theorem 4

If  $\Omega$  is a bounded simply-connected domain and if every boundary point of  $\Omega$  is simple, then every Riemann mapping of  $\Omega$  onto  $\mathbb{D}$  extends to a homeomorphism of  $\overline{\Omega}$  onto  $\overline{\mathbb{D}}$ .

## Proof.

Let  $f : \Omega \rightarrow \mathbb{D}$  be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend  $f$  to  $\overline{\Omega}$  using sequences. By 2, it follows that  $f$  so extended is one-one. We now check that it is continuous on  $\overline{\Omega}$ . As remarked earlier, this would finish the proof.

To this end, let  $(z_n)_n$  be an arbitrary sequence in  $\overline{\Omega}$  that converges to  $z$ . We can pick a corresponding sequence  $(\alpha_n)_n$  in  $\Omega$  such that  $|\alpha_n - z_n| < 1/n$  and  $|f(\alpha_n) - f(z_n)| < 1/n$ . Thus,  $\alpha_n \rightarrow z$  and hence,  $f(\alpha_n) \rightarrow f(z)$ . In turn,  $f(z_n) \rightarrow f(z)$ , as desired. □



# A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map  $S^1 \rightarrow \mathbb{C}$ .

## Corollary 5

If every boundary point of a bounded simply-connected region  $\Omega$  is simple, then the boundary of  $\Omega$  is a Jordan curve, and  $\overline{\Omega}$  is homeomorphic to  $\overline{\mathbb{D}}$ .

In fact, the converse is true too: If the boundary of  $\Omega$  is a Jordan curve, then every boundary point of  $\Omega$  is simple.

## Theorem 6 (Radial Limit Theorem)

To every  $g \in \mathcal{O}^\infty$  corresponds a function  $g^* \in L^\infty(S^1)$ , defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If  $g^*(e^{i\theta}) = 0$  for almost all  $e^{i\theta}$  on some arc  $J \subseteq S^1$ , then  $g(z) = 0$  for every  $z \in \mathbb{D}$ .

## Theorem 7 (Lindelöf's Theorem)

Suppose  $\Gamma$  is a curve such that  $\Gamma([0, 1)) \subseteq \mathbb{D}$  and  $\Gamma(1) = 1$ .

If  $g \in \mathcal{O}^\infty$  and

$$\lim_{t \rightarrow 1^-} g(\Gamma(t)) = L,$$

then  $g$  has radial limit  $L$  at 1.

# Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that  $f$  cannot be extended to  $\beta$ . Then, there exists a sequence  $(\alpha_n)_n$  in  $\Omega$  and points  $w_1, w_2 \in \overline{\mathbb{D}}$  such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose  $\gamma$  as given by  $\beta$  being a simple boundary point, and put  $\Gamma := f \circ \gamma$ . Let  $g = f^{-1}$  and put  $K_r := g(\overline{D}(0; r))$  for  $0 < r < 1$ . Then  $K_r$  is a compact subset of  $\Omega$ . Since  $\gamma(t) \rightarrow \beta$  as  $t \rightarrow 1$ , there exists  $t^* < 1$  (depending on  $r$ ) such that  $\gamma(t) \notin K_r$  if  $t^* < t < 1$ . Thus,  $|\Gamma(t)| \rightarrow 1$  as  $t \rightarrow 1$ . In particular,  $|w_1| = |w_2| = 1$ .

Let  $J$  be one of the open arcs of  $S^1 \setminus \{w_1, w_2\}$  such that every radius of  $\mathbb{D}$  which ends at a point of  $J$  intersects the range of  $\Gamma$  in a set which has a limit point on  $S^1$ . By the Radial Limit Theorem,  $g$  has radial limits a.e. on  $S^1$  since  $g \in \mathcal{O}^\infty$  (as  $\Omega$  is bounded).

## Continuing the Proof

We have that  $g \circ \Gamma = \gamma$  and that  $g$  has radial limits a.e. on  $S^1$ . Now, for whichever  $e^{it} \in J$  the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since  $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$  as  $t \rightarrow 1$ . Thus, by the Radial Limit Theorem again, applied to  $g - \beta$ , we see that  $g \equiv \beta$  on  $\mathbb{D}$ , contradicting that  $g$  is an injection.

Thus, we have shown that  $w_1 = w_2$  and  $|w_1| = 1$ .

## Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let  $\beta_1, \beta_2$  be simple boundary points with  $f(\beta_1) = f(\beta_2)$ . We may assume  $f(\beta_i) = 1$ . Let  $\gamma_i$  be curves with  $\gamma_i([0, 1)) \subseteq \Omega$ , and  $\gamma_i(1) = \beta_i$ , and let  $\Gamma_i := f \circ \gamma_i$ . Then, each  $\Gamma_i$  satisfies the condition of Lindelöf's Theorem with

$$\lim_{t \rightarrow 1} g(\Gamma_i(t)) = \lim_{t \rightarrow 1} \gamma_i(t) = \beta_i.$$

Thus, the radial limit of  $g$  at 1 is both  $\beta_1$  and  $\beta_2$  and hence,  $\beta_1 = \beta_2$ .  $\square$